



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Grazing Behaviour of Yak (*Bos grunniens*) in Warm-and Cold-Season Paddocks of *Potentilla fruticosa* Alpine Rangeland in Northern Qinghai-Tibetan Plateau

Nobumi Hasegawa
University of Miyazaki, Japan

R. Song
Yushu State Animal Husbandry and Veterinary Center, China

G. Li
Yushu Prefectural Grassland Station, China

Akira Fukuda
Shizuoka University, Japan

S. Feng
Mengyuan Prefectural Animal Husbandry and Veterinary Station, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/3-2/18>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Grazing behaviour of yak (*Bos grunniens*) in warm-and cold-season paddocks of *Potentilla fruticosa* alpine rangeland in Northern Qinghai-Tibetan Plateau

N. Hasegawa¹, R. Song², G. Li³, A. Fukuda⁴ and S. Feng⁵

¹ Faculty of Agriculture, University of Miyazaki, Gakuen-kibanadai, Miyazaki, 889-2192, Japan, E-mail: nhasegaw@cc.miyazaki-u.ac.jp, ² Yushu State Animal Husbandry and Veterinary Center, Jiegu Town, Yushu State, Qinghai Province, 815000, China, ³ Yushu Prefectural Grassland Station, Jiegu, Yushu State, Qinghai Province, 815000, China, ⁴ Graduate School of Science and Technology, Shizuoka University, Hamamatu, 432-8561, Japan, ⁵ Mengyuan Prefectural Animal Husbandry and Veterinary Station, Haomanzhen, Mengyuan Prefecture, Haibei State, Qinghai Province, 810300, China

Key words : yak, grazing behaviour, two-season-rotational grazing, alpine rangeland, Qinghai-Tibetan Plateau

Introduction Authors have been investigating the behaviour of yaks (*Bos grunniens*), botanical diversity and material circulation in rangelands of Qinghai-Tibetan Plateau (Hasegawa *et al.*, 2006; Song *et al.*, 2006). Two-season-rotational grazing system between warm-and cold-season paddocks is increasing in Qinghai-Tibetan Plateau under settlement policy of nomads by Chinese government. In this study, grazing behaviour of yak cows was investigated in northern rangeland of Qinghai-Tibetan Plateau to evaluate the two-season-rotational grazing system.

Materials and methods This study was carried out in August and December, 2005 in warm-and cold-season paddocks (WSP and CSP) of *Potentilla fruticosa* alpine rangeland in Mengyuan Prefecture, Haibei State, Qinghai Province, China. Behaviour of 3 yak cows was observed on the grazing time and bite numbers in a patch and the numbers of steps in a patch, between feeding stations and between patches. Positions of 5 yaks were recorded by GPS.

Results WSP was significantly greater than CSP ($p < 0.001$) in bite number per patch, bite rate in a patch, number of feeding station per patch and number of steps within a patch, and was significantly smaller than CSP ($p < 0.01$) in number of steps between patches (Table 1). Those two were not different in other items. Figure 1 shows the distribution of yaks in WSP and CSP. Distance of yaks from gateway of night paddock averaged 449.2 m in WSP and 334.2 m in CSP.

Table 1 Grazing behavior of yak cows in rotationally grazed warm- and cold-season paddocks in alpine rangeland.

Item	WSP ¹⁾	CSP ²⁾	p ³⁾
No. of patches observed	218	367	-
No. of patches visited, patches/min	0.9	1.0	ns
Grazing time, sec/patch	55.9	52.4	ns
Bite number, bites/patch	67.4	37.2	***
Bite rate within a patch, bites/sec	1.2	0.7	***
No. of feeding stations, FS/patch	7.1	5.9	***
No. of steps within a patch, steps/patch	6.1	2.5	***
Step rate within a patch, steps/sec	0.17	0.16	ns
Moving time between patches, sec	10.5	8.6	ns
No. of steps between patches, steps	4.4	5.8	**
Step rate between patches, steps/sec	0.86	0.85	ns

Values are expressed as mean. ¹⁾ WSP: paddock grazed in warm seasons and ²⁾ CSP: paddock grazed in cold seasons by yaks for over 20 years. ³⁾ * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and ns= no significance by Wilcoxon test.

Conclusions Behavioural difference of yaks between WSP and CSP was considered to show that yaks were not able to compensate the decreases of edible plants and intake even by changing grazing strategy in CSP.

Acknowledgment This research was supported by Grant-in-Aids for Scientific Research (A) No. 15255020 from Japan Society for Promotion of Science.

References

- Hasegawa N., Song R., Kozono M., Idota S., Nishiwaki A., Li G., Fukuda A., Zhou Q. (2006). Differences in yak (*Bos grunniens*) grazing behaviour and chemical composition of feces in the southern and northern Qinghai-Tibetan Plateau in China. *Acta Prataculturae Sinica*. 15 (suppl.). 286-288.
- Song R., Hasegawa N., Idota S., Li G., Nishiwaki A., Jiu C., Xu N., Zhou Q. (2006). Botanical composition, aboveground biomass and grazing behaviour of yak (*Bos grunniens*) in the southern rangeland of Qinghai Province, China. *Acta Prataculturae Sinica*. 15 (suppl.). 289-291.




Figure 1 Distribution of yaks in WSP¹⁾ and CSP²⁾ by the distance from gateway of night paddock. ^{1 2)} Refer to Table 1.